

First In the world

Beyond the Dream



Hankook Korus Pharm. Co., Ltd.,

Hankook Korus Pharm. Co., Ltd., as a leading pharmaceutical company that loves life and contributes to human society,
will grow into a global bio company loved and trusted by customers.

Chuncheon Bio Plant received KGMP certification in 2019 for biopharmaceuticals and produces biopharmaceuticals such as PEG interferon alpha-2b (PEG-IFN alpha-2b), Erythropoietin (EPO), and Human Growth Hormone (rhGH). It has a one-stop line from bio stock solution to charging, and bio-finished products can be produced up to 18 million vials and PFS per year.

Main Production equipment

Bio Drug Substance (DS)

Cultivation Line (Upstream)

: 15L, 20L, 50L, 100L, 130L, 200L, 1,000L

Purification Line (Downstream)

: Depth Filtration system, TFF(UF/DF)system, Chromatography system

Drug Product (DP)

Prefilled Syringe Line

- Syringe Filling & Closing
- Load assembly & Labeling Machine
- Blister Machine

Vial Line

- Vial Washing
- Fill & Stoppering
- AI-Cap sealing
- Tunnel sterilizer
- Complete line

Freeze Dryer

Acyclor Inj. 250mg

Antiviral medication

Active Ingredient

Acyclovir 250mg / mL

Indications

- Herpes Simplex Infections in Immunocompromised Patients: Aciclovir for injection is indicated for the treatment of initial and recurrent mucosal and cutaneous herpes simplex (HSV-1 and HSV-2) in immunocompromised patients.
- Initial Episodes of Herpes Genitalis: Aciclovir for injection is indicated for the treatment of severe initial clinical episodes of herpes genitalis in immunocompetent patients.
- Herpes Simplex Encephalitis: Aciclovir for injection is indicated for the treatment of herpes simplex encephalitis.
- Varicella-Zoster Infections in Immunocompromised Patients: Aciclovir for injection is indicated for the treatment of varicella-zoster (shingles) infections in immunocompromised patients.

Dosage and administration

Injections are slowly instilled intravenously over 1 hour or more each time it is administered. The duration of administration is appropriate for 5 days, but it can be adjusted according to the patient's symptoms and healing status. Herpes simplex encephalitis is usually administered for 10 days.

new product



Zinc Power Inj. 10mL

Mineral Supplement

Active Ingredient

Zinc Sulfate Hydrate 43.9 mg / 10mL (As Zinc 10 mg)

Indications

Zinc supplementation in Total Parenteral Nutrition (TPN)

Dosage and administration

Division	Zinc Permissible Dosage Range
Metabolically stable TPN patients	2.5-4 mg/day
pediatric patient	Under 5 years old: 100 ug/kg/day Premature infant(3 kg or less): 300 ug/kg/day
Adult patients with fluid loss in the small intestine	Additional 12.2 mg/L (fluid loss) or additional 17.1 mg/kg (excretion)
acute catabolism	4.5-6mg/day

new product



Kotamin C Inj. 500mg

Vitamin C supplement

Active Ingredient

Ascorbic acid 500 mg / mL

Indications

1. Prevention and treatment of scurvy.
2. Vitamin C supplementation.
3. Vitamin C deficiency

Dosage and administration

As ascorbic acid, 50 to 1,000 mg per day, divided once to several times and injected subcutaneously, intramuscularly or intravenously. Increase or decrease appropriately according to age and symptoms.

new product



API Supply Available

DMF

Hyaldase Inj. / MaxHyal Inj.

Enzyme, Disease of the blood and hematopoietic organs

Active Ingredient

1. Hyaldase Injection (Low Purity Hyaluronidase) : Hyaluronidase 1,500IU
2. MaxHyal Injection (High Purity Hyaluronidase) : Hyaluronidase 1,500IU

Indications

- Increased penetration in case of subcutaneous or intramuscular Injection, local anesthetic and subcutaneous injection
- Promote reabsorption of body fluids and blood that are excessively present in tissues

Dosage and administration

1. Hyaldase Injection : Liquide (Vial)
2. MaxHyal Injection : Lyophilized powder (Vial)

new product



Developing Product List

No	Product Name	Active Ingredient	Product Type	Note
01	Vancomycin Inj. 0.5g	Vancomycin 0.5g	Vial (Lyophilized)	
02	Vancomycin Inj. 1.0g	Vancomycin 1.0g	Vial (Lyophilized)	
03	Kortamin C Inj. (10g/20mL)	Ascorbic Acid 10g/20mL	Vial (Solution)	
04	Low Purity Hyaladase 1,500IU Inj.	Hyaluronidase 1,500IU	Vial (Lyophilized)	
05	Ibuprofen Inj.	Ibuprofen 400mg/4mL	Vial (Solution)	
06	High Purity Hyaladase 1,500IU Inj.	Hyaluronidase 1,500IU	Vial (Solution)	
07	Ibaron Inj.	Ibandronate Acid 3mg	PFS	
08	Ontron Inj.	Ondansetron 4mg	Vial (Solution)	
09	Nekopam HCl 40mg Inj.	Nefopam HCl 40mg	Vial (Solution)	
10	Panzole 40mg Inj.	Pantoprazole 40mg	Vial (Lyophilized)	
11	Teplanin 200mg Inj.	Teicoplanin 200mg	Vial (Lyophilized)	
12	Elbrakan Inj.	Fulvestrant 50mg	PFS	
13	Amika Inj.	Amikacin Sulfate 500mg	Vial (Solution)	
14	Immusin Inj. (Thymosin α 1)	Thymosin α 1 1.6mg	Vial (Lyophilized)	
15	Polydyron Inj. (PDRN)	PDRN 5.625mg	Vial (Solution)	
16	Netisul Inj. (Netylmycin Sulfate)	Netylmycin Sulfate 150mg	Vial (Solution)	
17	Amodar Inj. (Amiodaron)	Amiodaron 150mg	Vial (Solution)	
18	Tobracin Inj.	Tobramycin 80mg	Vial (Solution)	
19	Gluthathion Inj.	Gluthathion 1,200mg	Vial (Lyophilized)	
20	Hydroxocobalamin Inj. 2.5mg	Hydroxocobalamin 2.5mg	Vial (Solution)	
21	Sodium Heparin 25,000IU Inj.	Sodium Heparin 25,000IU	PFS	
22	Enoxaparin 80mg PFS Inj.	Enoxaparin 80mg	PFS	
23	Hydroxocobalamin Inj. 5mg	Hydroxocobalamin 5mg	Vial (Solution)	
24	Cerebrolysin Inj. 212.5mg	Cerebrolysin 212.5mg	Vial (Solution)	
25	Ferinject inj 10mL	Iron 500mg/10ml	Vial (Solution)	
26	Human placenta ext. 1mL Inj.	Human placenta ext. 1mL	Vial (Solution)	
27	Sugammadex sodium Inj.	Sodium Sugammadex 100mg	Vial (Solution)	

Rebistar Tab.

Drug for peptic ulcer and gastro-oesophageal reflux disease

Active Ingredient

Rebamipide 100mg

Indications

- Gastric ulcer
- Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in the following diseases: acute gastritis, acute exacerbation of chronic gastritis

Dosage and administration

Adult: 100 mg of Rebistar Tab. is orally administered 3 times a day.
However, in the case of gastric ulcer, it should be administered before breakfast, evening and bedtime.

Alimentary Tract & Metabolism



Udica Tab.

Secondary bile acid

Active Ingredient

Ursodeoxycholic Acid 300mg, 500mg

Indications

Improvement of Liver function in Primary biliary cirrhosis (PBC)

Dosage and administration

Adult: Administer 300mg at a time, 3times a day during meal

Alimentary Tract & Metabolism



Dyryl Tab.

2nd Generation sulfonylurea antidiabetic drug

Active Ingredient

Glimepiride 2mg

Indications

Dyryl Tab. is indicated for the treatment of type 2 diabetes mellitus, when diet, physical exercise and weight reduction alone are not adequate.

Dosage and administration

The dosage is determined by the results of blood and urinary glucose determinations.
The starting dose is 1 mg glimepiride per day. The maximum recommended dose is 6 mg glimepiride per day.

Alimentary Tract & Metabolism



Nexome Tab. 20mg/40mg

Treatment of GERD & ZES

Active Ingredient

Nexome Tab. 20mg
(a)Esomeprazole Magnesium Dihydrate 21.69 mg(as esomeprazole 20mg)
Nexome Tab. 40mg
(b)Esomeprazole Magnesium Dihydrate 43.38 mg(as esomeprazole 20mg)

Indications




















- Gastroesophageal reflux disease
- Antibiotic combination therapy for Helicobacter pylori eradication
- Short-term therapy for upper gastrointestinal symptoms associated with non-steroidal anti-inflammatory analgesics
- Patients who need continuous nonsteroidal anti-inflammatory medication
- Treatment of Zollinger-Ellison syndrome
- Maintenance therapy after induction of re-hemorrhage by stomach ulcer or duodenal ulcer by intravenous injection















Dosage and administration

For adults, administered 20mg a time and once a day.
In the treatment of erosive reflux disease, treatment with Zollinger-Ellison syndrome, maintenance therapy, administered 40mg a time and once a day.

Alimentary Tract & Metabolism



Product Name (Active Ingredients & Strength)		Indications	Remark
Almagent Tab. Almagate 500mg		Gastric or duodenal ulcer, Gastritis, Hypergastric acidity (Heartburn, Nausea, Vomiting, Gastralgia, Sour belching), Reflux esophagitis	
DDB Cap. Biphenyl Dimethyl Dicarboxylate Pills (7.5mg as Biphenyl Dimethyl Dicarboxylate) 75mg		Chronic persistent and active hepatitis B, Liver cirrhosis, Drug-induced hepatitis associated with high SGPT level	
DDB Plus Cap. Biphenyl Dimethyl Dicarboxylate 12.5mg Ursodeoxycholic acid 50mg		Chronic persistent hepatitis with elevated SGPT	
Domperidone-K Tab. Domperidone Maleate (10mg as Domperidone) 12.7mg		The relief of the symptoms of nausea and vomiting, Epigastric sense of fullness, Upper abdominal discomfort and regurgitation of gastric contents	
Famostop Tab. Famotidine 40mg		Active duodenal ulcer, Duodenal ulcer, Active benign gastric ulcer, Gastroesophageal reflux disease (GERD), Pathological hypersecretory conditions	
Korus Ascorbic Acid Tab. Coated Ascorbic Acid (500mg as Ascorbic Acid) .. 515.5mg		Scurvy, Decreased strength Physical fatigue, Capillary bleeding, Pigmentation during period of lactation and illness	
Korus Cimetidine Tab. Cimetidine 400mg		Active duodenal ulcer, active benign gastric ulcer, Erosive gastroesophageal reflux disease (GERD), Prevention of upper gastrointestinal bleeding, pathological hypersecretory conditions	
Korus Omeprazole Cap. (a)Omeprazole Granule 115mg (b)Omeprazole Granule 233mg		Duodenal ulcer, Gastritis, Gastroesophageal reflux disease (GERD), Erosive Esophagitis, Pathological Hypersecretory Conditions	
Rabepa Tab. (a)Rabeprazole sodium 10mg (b)Rabeprazole sodium 20mg		Erosive or ulcerative GERD, Symptomatic GERD, Duodenal ulcers, Helicobacter pylori eradication to reduce the risk of duodenal ulcer recurrence, Pathological hypersecretory conditions including Zollinger-Ellison Syndrome	
Ranitidine Tab. (a)Ranitidine Hydrochloride (75mg as Ranitidine) 84mg (b)Ranitidine Hydrochloride (300mg as Ranitidine) 336mg		Active duodenal ulcer, Pathological hypersecretory conditions, Gastric ulcer, Erosive gastroesophageal reflux disease (GERD), Endoscopically diagnosed erosive esophagitis	
UDCA Tab. (a)Urosodeoxycholic Acid 200mg (b)Urosodeoxycholic Acid 300mg		Liver disease due to bile secretion deficiency Biliary tract disease, Chronic liver disease, Dyspepsia Cholelithiasis, Primary Biliary Cirrhosis(PBC)	
Zanide Cap. Nizatidine 150mg		Duodenal ulcer, Benign gastric ulcer, Prevention of duodenal or benign gastric ulcer recurrence, Gastric oesophageal reflux disease (including erosions, ulcerations and associated heartburn), Gastric and/or duodenal ulcer associated with concomitant use of non-steroidal anti-inflammatory drugs	
Max D's Tab. Calcium carbonate 750mg Cholecalciferol Concentrate Powder 10mg		Prevalence of vitamin D: Pregnancy, lactation, development, old age Calcium deficiency and other calcium supply	
Udica Tab. Ursodeoxycholic acid 500mg		Liver disease due to bile secretion deficiency Biliary tract disease, Chronic liver disease, Dyspepsia Cholelithiasis, Primary Biliary Cirrhosis(PBC)	
Itopam Tab. Itopride hydrochloride 50mg		Functional Digestive symptoms due to indigestion (abdominal distension, abdominal pain, anorexia, heartburn, nausea, vomiting)	
Dyryl Tab. Glimepiride 2mg		Dyryl Tab. is indicated for the treatment of type 2 diabetes mellitus, when diet, physical exercise and weight reduction alone are not adequate.	
A-formin SR Tab. Metformin Hydrochloride 500mg		A-formin SR Tab. is indicated for the treatment of type 2 diabetes mellitus, when diet, physical exercise and weight reduction alone are not adequate (In particular, overweight diabetic patients).	
Piota Tab. Pioglitazone Hydrochlorid .. 16.53mg		Piota Tab. is indicated for the treatment of type 2 diabetes mellitus, when diet, physical exercise and weight reduction alone are not adequate.	
Dyryl-M Tab. Glimepiride 2mg Metformin Hydrochloride 500mg		In combination with dietary and exercise therapy in patients with non-insulin-dependent diabetes mellitus (type 2) - Administered when there is insufficient blood glucose control effect by Glimepiride or Metformin monotherapy. - Replacement of Glimepiride and Metformin combination therapy	

Product Name (Active Ingredients & Strength)		Indications	Remark
Rebistar Tab. Rebamipide 100mg		Gastric ulcer Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema): Acute gastritis, acute exacerbation of chronic gastritis	
Spalen Tab. Artemisia Herb 95% Ethanol Soft Ext. (20→1)..... 60mg		Improve mucosal lesions (erosion, bleeding, redness, edema): acute gastritis, chronic gastritis Prevention of gastritis by administration of non-steroidal anti-inflammatory drugs(NSAID)	
Nexome Tab. (a)Esomeprazole Magnesium Dihydrate ... 21.69 mg(as esomeprazole 20mg) (b)Esomeprazole Magnesium Dihydrate ... 43.38 mg(as esomeprazole 40mg)	 	Gastroesophageal reflux disease, Antibiotic combination therapy for Helicobacter pylori eradication, Short-term therapy for upper gastrointestinal symptoms associated with non-steroidal anti-inflammatory analgesics, Patients who need continuous nonsteroidal anti-inflammatory medication, Treatment of Zollinger-Ellison Syndrome, Maintenance therapy after induction of re-hemorrhage by stomach ulcer or duodenal ulcer by intravenous injection	
Korus Levosulpiride Tab. Levosulpiride 25mg		Relief of the following symptoms due to functional dyspepsia: Abdominal bloating, upper abdominal discomfort, heartburn, trim, nausea, vomiting	
Korus Tiropramide HCl Tab. Tiropramide HCl 100mg		Acute spastic pain in the following diseases: Gastrointestinal colic, abdominal colic due to various causes, colic of the kidneys and uretersAbdominal cramps and pain in the following diseases: Gastrointestinal motility disorder, gallstone disease, cholecystitis, postoperative adhesions	
Ocpaminin Tab. Octylonium Bromide 20mg		Gastrointestinal cramps and irritable bowel syndrome	
Korus Mosapride Tab. Mosapride Citrate Hydrate ... 5.29mg		Functional Digestive symptoms due to indigestion (heartburn, nausea, vomiting)	
Trimetine Tab. Trimebutine Maleate 150mg		Esophageal reflux, hiatus hernia, dyspepsia (abdominal pain, indigestion, nausea, vomiting) Irritable bowel syndrome and spastic colon Pediatric disease: addictive vomiting, non-infectious intestinal tract obstruction (constipation, diarrhea), agitation, sleep disorder	
Octic Acid HR 600mg Thioctic Acid 600mg		Mitigation of diabetic polyneuritis	
Aroct Acid Tab. 480mg R-Thioctic Acid Tromethamine 480mg		Mitigation of diabetic polyneuritis	
Piota-M Tab. Metformin Hydrochloride 480mg Pioglitazone Hydrochloride .. 16.53mg		Metformin or pioglitazone alone can not control enough blood sugar Replacement therapy with pioglitazone and metformin	
Elbis Tab. Ranitidine Hydrochloride 84mg Sucralfate Hydrate 300mg Tripotassium Bismuth Dicitrate .. 100mg		Gastrointestinal / duodenal ulcer due to gastric and duodenal ulcers, gastritis, Zollinger-Ellison syndrome, reflux esophagitis, pre-narcotic medication (prevention of Mendelson syndrome), postoperative ulcers, nonsteroidal anti-inflammatory drugs(NSAID)	
Bearase Tab. Pancreatin 175mg Dimethicone 175mg Hemicellulase 175mg Ox bile extract 175mg		Dyspepsia, loss of appetite (loss of appetite), overeating, impairment, promotion of digestion,	

K-Pirome Inj. 0.5g/1.0g

4th Generation Cephalosporin Antibiotic

Active Ingredient

Cefpirome Sulfate 0.5g/1.0g

Indications

Infectious disease caused by Bacteria That Susceptibility to Cefpirome

- Upper and lower urinary tract infections with complications
- Lower respiratory tract infections
- Skin and soft tissue infections
- Infectious disease when neutrophil reduction Sepsis

Dosage and administration

- Upper and Lower Urinary Tract Infections with Complications
 - : 1dose 1g / Interval 12 hours / Total 2g
- Skin and Soft Tissue Infectious Diseases
 - : 1dose 1g / Interval 12 hours / Total 2g
- Lower respiratory tract infections
 - : 1dose 1g or 2g / Interval 12 hours / Total 2g or 4g
- blood poisoning
 - : 1dose 2g / Interval 12 hours / Total 4g
- Infections When Neutrophil Reduction
 - : 1dose 2g / Interval 12 hours / Total 4g

Anti-infectives for Systemic Use



Cepim Inj.

4th Generation Cephalosporin Antibiotic

Active Ingredient

Cefepime Hydrochloride 1g, 2g (potency)

Indications

Pneumonia, Empiric therapy for febrile neutropenia, Uncomplicated and complicated Urinary Tract Infections
Complicated Intra-abdominal Infections (used in combination with metronidazole)

Dosage and administration

1. Adults and children over 12 years old:
According to the symptoms, 1 ~ 2g (potency) per day is divided into 2, intravenous injection or intramuscular injection.
2. Children from 2 months to 12 years
Recommended dose is 50 mg / kg body weight and is injected intravenously every 8 hours 3 times a day

Anti-infectives for Systemic Use



Korus Ceftriaxone Inj.

3rd generation cephalosporin product

Active Ingredient

Ceftriaxone sodium 0.25g, 0.5g, 1g, 2g (potency)

Indications

Pneumonia, Septicaemia, Meningitis, Skin and soft tissue infections,
Infections in neutropenic patients, Gonorrhoea, Peri-operative prophylaxis of infections associated with surgery.

Dosage and administration

Adults and children 12 years and over
Standard therapeutic dosage: 1~2g (potency) once daily.
Children under 12 years
Standard therapeutic dosage: 20-50mg (potency) /kg body-weight once daily.
Do not exceed 50mg (potency) per Kg of body weight.

Anti-infectives for Systemic Use



Korus Cefotiam Inj.

2th Generation Cephalosporin

Active Ingredient

Cefotiam Hydrochloride 0.5g, 1g (potency)

Indications

Susceptible strains

Staphylococcus, Streptococcus(excluding Enterococci), Streptococcus pneumoniae, Haemophilus influenzae, Escherichia coli, Klebsiella, Enterobacter sp., Citrobacter, Proteus mirabilis, Proteus vulgaris, Proteus rettgeri and Proteus morganii.

Major indications

Septicemia Infections to a burn or operative wound, subcutaneous abscess, carbuncle, furuncle, furunculosis, osteomyelitis, pyogenic arthritis, tonsillitis (peritonsillitis, peritonsillar abscess), bronchitis, bronchiectasis with infections, pneumonia, pulmonary suppuration, pyothorax, cholangitis, cholecystitis, peritonitis, mephropyelitis, cystitis, urethritis, prostatitis, meningitis, intrauterine infections, pelvic inflammatory disease, parametritis, adnexitis, bartholinitis, otitis media and sinusitis

Dosage and administration

I.V. Injection

Adult : Administer 0.5 to 2 g (potency) of cefotiam hydrochloride in 2 to 4 times divided doses a day by I.V.

Children : Administer 40 to 80 mg (potency) of cefotiam hydrochloride per kilogram in 3 to 4 times divided doses a day by I.V.

The dose for adult's septicemia can be increased up to 4 grams (potency) a day.

The dose for child can be increased up to 160 mg (potency)/kg/day with serious or refractory infections, such as septicemia

and meningitis. The dosage may be adjusted according to the patient's age and symptom.

I.M. Injection

Adult : Administer 0.5 to 2 g (potency) of cefotiam hydrochloride in 2 to 4 times divided doses a day by I.M. The dosage may be adjusted according to the patient's age and symptom.

Use after dissolving cefotiam for injection in 3 mL of solvent which is attached 0.5% lidocaine per vial.

Anti-infectives for Systemic Use



Cefaclor Cap. 250mg

2th Generation Cephalosporin

Active Ingredient

Cefaclor 250mg(potency)

Indications

Cefaclor is indicated in the treatment of the following infections :

- Otitis media infection
- Lower respiratory infections including pneumonia
- Pharyngitis and tonsillitis
- Urinary tract infections including pyelonephritis and cystitis
- skin and skin structure infections

Dosage and administration

Adults : The usual adult dosage is 250 mg every 8 hours.

For more severe infections (such as pneumonia) or those caused by less susceptible organisms, doses may be doubled. Cefaclor may be administered in the presence of impaired renal function.

Under such a condition, the dosage usually is unchanged.

In the treatment of p-hemolytic streptococcal infections, a therapeutic dosage of Cefaclor should be administered for at least 10 days.

Anti-infectives for Systemic Use



Flunazole Cap.

1st triazole antifungal medication

Active Ingredient

Fluconazole 50mg, 150mg

Indications

Cryptococcal meningitis, Coccidioidomycosis, Invasive candidiasis., Mucosal candidiasis including oropharyngeal, oesophageal candidiasis, candiduria and chronic mucocutaneous candidiasis, Chronic oral atrophic candidiasis (denture sore mouth) if dental hygiene or topical treatment are insufficient, Vaginal candidiasis, acute or recurrent; when local therapy is not appropriate, Candidal balanitis when local therapy is not appropriate, Dermatomycosis including tinea pedis, tinea corporis, tinea cruris, tinea versicolor and dermal candida infections when systemic therapy is indicated, Tinea unguinum (onychomycosis) when other agents are not considered appropriate.

Dosage and administration

In Adults aged 16 – 60 years:

Vaginal candidiasis or candidal balanitis – 150 mg single oral dose.

Anti-infectives for Systemic Use



Tobracin Inj. 80mg

Aminoglycoside antibacterial

Active Ingredient

Tobramycin 80 mg / 2mL

Indications

Tobramycin Injection is indicated in the treatment of the following serious infections caused by susceptible micro-organisms:

- The treatment of central nervous system infections including meningitis, septicaemia and neonatal sepsis;
- The treatment of gastro-intestinal infections including peritonitis;
- The treatment of complicated and recurrent urinary tract infections such as pyelonephritis and cystitis;
- The treatment of lower respiratory tract infections, including pneumonia, bronchopneumonia and acute bronchitis;
- The treatment of skin, bone and soft tissue infections including burns

Dosage and administration




















For intramuscular injection, dissolve in 1.5mL of physiological saline injection and water for injection.

















For intravenous injection, dilute in 50-100 mL of physiological saline or 5% glucose injection and inject intravenously for 20-60 minutes.












In case of severe infection, tobramycin is administered intravenously or intramuscularly by dividing 3 mg per kg of body weight per day into 3 divided doses.

Anti-infectives for Systemic Use



Product Name (Active Ingredients & Strength)	Indications	Remark
Korus Cefotaxime Sodium Inj. Cefotaxime Sodium 250mg/500mg/1g/2g 	Lower respiratory tract infections, Genitourinary infections, Gynecologic infections, Bacteremia/Septicemia, Skin and skin structure infections, Intra-abdominal infections, Bone and/or joint infections, Central nervous system infections	
Korus Cefotiam Hydrochloride Inj. Cefotiam Hydrochloride-Dried Sodium Carbonat 0.5g/1g 	Pneumonia, Bronchitis, Biliary tract infections, Peritonitis, Urinary tract infections, Infections caused by surgery, Trauma and sepsis.	
Korus Ceftriaxone Sodium Inj. Ceftriaxone Sodium 250mg/500mg/1g/2g 	Pneumonia, Septicaemia, Meningitis, Skin and soft tissue infections, Infections in neutropenic patients, Gonorrhoea, Perioperative prophylaxis of infections associated with surgery	
Korus Cefuroxime Sodium Inj. (a)Cefuroxime Sodium 250mg (b)Cefuroxime Sodium 750mg (a)Cefuroxime Sodium 1.5g 	Acute and chronic bronchitis, Infected bronchiectasis, Bacterial pneumonia, Lung abscess and postoperative chest infections, Sinusitis, Tonsillitis and pharyngitis, Acute and chronic pyelonephritis, Cystitis and asymptomatic bacteriuria, Cellulitis	
Korus Cycloserine Cap. Cycloserine 250mg 	Active pulmonary and extra-pulmonary tuberculosis, Acute urinary tract infections	
Korus Doxycycline Cap. Doxycycline Hyclate Hydrate .. 100mg 	Pneumonia and other lower respiratory tract infections, Urinary tract infections, Infections due to Chlamydia trachomatis	
Korus Ethambutol Hydrochloride Tab. Ethambutol Hydrochloride .. 200mg/400g 	Pulmonary tuberculosis, Other tuberculosis	
Korus Fluconazole Cap. Fluconazole 50mg/150mg 	Vaginal candidiasis, Acute or recurrent or candidal balanitis associated with vaginal candidiasis	
Korus Ofloxacin Tab. Ofloxacin 100mg/200mg/400mg 	Urinary tract infection, Respiratory tract infections, Uncomplicated urethral and cervical gonorrhea	
Korus Rifampicin Tab. Rifampicin 450mg/600mg 	Tuberculosis, Leprosy, Brucellosis, Legionnaires Disease and serious staphylococcal infections, Meningococcal meningitis, Haemophilus influenzae	
Korus Rifampicin Cap. Rifampicin 150mg/300mg 		
Levotacin Tab. Levofloxacin .. 100mg/250mg/500mg 	Acute bacterial sinusitis, Acute exacerbations of chronic bronchitis, Community-acquired pneumonia, Complicated skin and soft tissue infections, Pyelonephritis and complicated urinary tract infections, Chronic bacterial prostatitis	
Cefaxin Cap. Cephalexin Hydra 250mg/500mg 	Respiratory tract infections, Urinary tract infections, Skin and soft tissue infections, Otitis media	
Mepenem Inj. Meropenem Dried Sodium Carbonate 604mg/1208mg 	Pneumonia, Broncho-pulmonary infections, Complicated urinary tract infections, Complicated intra-abdominal infections	
Metronidazole Tab. Metronidazole 250mg/500mg 	Postoperative infections due to anaerobic bacteria, Septicaemia, Bacteraemia, Peritonitis, Brain abscess, Necrotising pneumonia,	
Sulfozone Inj. Sulbactam Sodium m · Cefoperazone Sodium 500mg/ 1g/ 2g 	Respiratory tract infections, Urinary tract infections, Peritonitis, Cholecystitis, Cholangitis and other intra-abdominal infections, Bacterial sinusitis, Meningitis, Skin and soft tissue infections	
Tetozole Tab. Ketoconazole 200mg 	Systemic fungal infections	
Latamox Inj. Latamoxef Sodium 0.25g/ 0.5g/1g 	Bone and joint infection, GI infections, Gynecological infections, Meningitis, Respiratory tract infections, Septicaemia, Skin infections	
K-flu Cap. 30mg/45mg/75mg (a) Oseltamivir Phosphate ... 39.4mg (b) Oseltamivir Phosphate ... 59.1mg (c) Oseltamivir Phosphate ... 98.5mg 	Treatment of influenza Prevention of influenza	

Product Name (Active Ingredients & Strength)		Indications	Remark
Floxam Inj. Flomoxef Sodium 0.5g/ 1.0g		Septicemia, Infectious endocarditis, Secondary infections in trauma, Burns, Surgical wounds, Pharyngolaryngitis, Tonsillitis, Secondary infections in chronic respiratory disease	
K-Mentine Tab. (A) Amoxicillin Hydrate 250mg Potassium clavulanate .. 212.73mg (B) Amoxicillin Hydrate 500mg Potassium clavulanate .. 212.73mg		Chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, Septic abortion	
Famcle Tab. Famciclovir 250mg/750mg		1. Shingles virus infection 2. Treatment of genital herpes infection and inhibition of recurrent genital herpes	
Korus Moxifloxacin Tab. Moxifloxacin Hydrochloride .. 400mg		1) Respiratory infections: Acute sinusitis, acute exacerbation of chronic bronchitis, pneumonia 2) Skin and soft tissue infection 3) Merged intraabdominal infection 4) Simple pelvic infection	
Cefurosaxime Tab. Cefuroxime axetil 250mg		Cefuroxime is a broad-spectrum bactericidal cephalosporin antibiotic that is very stable to beta-lactamase and also works against most ampicillin and amoxicillin-resistant strains.	
Korus Cefaclor Cap. (a) Cefaclor Hydrate 250mg (b) Cefaclor Hydrate 500mg		Pharyngitis and tonsillitis and as part of the management of Pneumonia, Bronchitis, Exacerbations of chronic bronchitis, sinusitis, Otitis media, Skin and soft tissue infections, Pyelonephritis and cystitis, Acute and chronic urinary tract infections	
Cefazol Inj. (a) Cefazolin Sodium 0.5g (b) Cefazolin Sodium 1g (c) Cefazolin Sodium 2g		Respiratory tract infections, Urinary tract infections, Biliary tract infections, Bone and joint infections, Genital infections, Septicemia, Endocarditis, Perioperative prophylaxis	
Cepim Inj. (a) Cefepime Hydrochloride 1g (b) Cefepime Hydrochloride 2g		Pneumonia, Empiric therapy for febrile neutropenic patients, Uncomplicated and complicated urinary tract infections, Uncomplicated skin and skin structure infections, Complicated intra-abdominal infections	
Ciploxacin Tab. (a) Ciprofloxacin Hydrochloride (250mg as Ciprofloxacin) 291mg (b) Ciprofloxacin Hydrochloride (500mg as Ciprofloxacin) 582mg		Urinary tract infections, Acute uncomplicated cystitis in females, Chronic bacterial prostatitis, Lower respiratory tract infections, Acute sinusitis, Skin and skin structure infections, Bone and joint infections, Complicated intra-abdominal infections, Infectious diarrhea, Typhoid fever, Uncomplicated cervical and urethral gonorrhea, Pediatric patients, Complicated urinary tract infections, Pyelonephritis, Inhalational Anthrax	
Clacin Tab. (a) Clarithromycin 250mg (b) Clarithromycin 500mg		Bacterial pharyngitis, Mild to moderate community-acquired pneumonia, Acute bacterial sinusitis, Acute exacerbation of chronic bronchitis, Skin and soft tissue infections of mild to moderate severity.	
Clicin Cap. (a) Clindamycin Hydrochloride .. 150mg (b) Clindamycin Hydrochloride .. 300mg		Severe infections caused by susceptible gram-positive aerobic organisms, staphylococci, streptococci and pneumococci or by susceptible anaerobic organisms.	
Fosfocin Inj. Fosfomicin Sodium 1g		Respiratory tract infections, Skin and soft tissue infections, Intestinal tract infections, Urinary tract infections, Blood poisoning, Peritonitis, Meningitis, Osteomyelitis, Adnexitis, Intra-uterine infection and pelvic inflammation	
Keradin Cap. (a) Cefradine Hydrate 250mg (b) Cefradine Hydrate 500mg		Sinusitis, Pharyngitis, Tonsillitis, Laryngo-tracheo bronchitis and otitis media, Acute and chronic bronchitis, Lobar and bronchopneumonia, Cystitis, Urethritis, Pyelonephritis, Impetigo, Abscess, Cellulitis, Furunculosis	
Kocezone Inj. (a) Ceftizoxime Sodium 0.5g (b) Ceftizoxime Sodium 1g		Lower respiratory tract infections, Urinary tract infections, Gonorrhoea, Pelvic inflammatory disease, Intra-abdominal infections, Septicemia, Skin and skin structure infections, Bone and joint infections, Meningitis	
Korazon Inj. (a) Cefoperazone Sodium 0.5g (b) Cefoperazone Sodium 1g (c) Cefoperazone Sodium 1g		Respiratory tract infections, Peritonitis and other intra-abdominal infections, Bacterial septicemia, Infections of the skin and skin structures, Pelvic inflammatory disease, Endometritis, and other infections of the female genital tract, Urinary tract infections, Enterococcal infections	
Korudim Inj. (a) Ceftazidime Hydrate - Dried Sodium Carbonate 0.5g (b) Ceftazidime Hydrate - Dried Sodium Carbonate 1g (c) Ceftazidime Hydrate - Dried Sodium Carbonate 2g		Nosocomial pneumonia, Broncho-pulmonary infections in cystic fibrosis, Bacterial meningitis, Chronic suppurative otitis media, Malignant otitis externa, Complicated urinary tract infections, Complicated skin and soft tissue infections, Complicated intra-abdominal infections, Bone and joint infections, Peritonitis associated with dialysis in patients on CAPD.	

Product Name (Active Ingredients & Strength)	Indications	Remark
Korus Pyrazinamide Tab. (a) Pyrazinamide 250mg (b) Pyrazinamide 500mg 	Active tuberculosis caused by Mycobacterium tuberculosis	
K-Pirom Inj. (a) Cefpirome Sulfate 0.5g (b) Cefpirome Sulfate 1.0g 	Infections caused by Cefpirome-sensitive organisms including lower respiratory tract infections, Complicated upper and lower urinary tract infections, Bacteraemia, Septicaemia, Infections in neutropenic and immunocompromised patients, Skin and soft tissue infections, Severe infections in intensive care patients.	
Panzedin Inj. Cefradine Hydrate 1g 	Sinusitis, Pharyngitis, Tonsillitis, Laryngo-tracheo bronchitis and otitis media, Acute and chronic bronchitis, Lobar and bronchopneumonia, Cystitis, Urethritis, Pyelonephritis, Impetigo, Abscess, Cellulitis, Furunculosis	
Cefacil Cap. Cefadroxil Monohydrat 500mg 	Pyelonephritis, cystitis, urethritis, skin and skin tissue infection, pharyngitis, tonsillitis	
Korus Acyclovir Tab. Acyclovir 200mg 	Treatment and prevention of herpes simplex virus infections of skin and mucosal tissues including acute and recurrent genital herpes. It is effective for the treatment of herpes zoster virus infection, especially acute pain. On the other hand, the effect on postherpetic neuralgia has not been proved yet. Varicella treatment in children over 2 years	
Itanox Tab. Itraconazole 100mg 	Cefuroxime is a broad-spectrum bactericidal cephalosporin antibiotic that is very stable to beta-lactamase and also works against most ampicillin and amoxicillin-resistant strains	
K-Levo Inj. Levofloxacin 5.1246mg/mL 	Community-acquired pneumonia, Complicated skin and soft tissue infections, For the above-mentioned infections levofloxacin should be used only when it is considered inappropriate to use antibacterial agents that are commonly recommended for the initial treatment of these infections, Pyelonephritis and complicated urinary tract infections, Chronic bacterial prostatitis, Inhalation anthrax: postexposure prophylaxis and curative treatment, Consideration should be given to official guidance on the appropriate use of antibacterial agents.	
K-pro Inj. Ciprofloxacin 2mg/mL 	<ul style="list-style-type: none"> · Lower respiratory tract · Chronic suppurative otitis media · Acute exacerbation of chronic sinusitis especially if these are caused by Gram-negative bacteria · Urinary tract infections · Genital tract infections · Infections of the gastro-intestinal tract (e.g. travellers diarrhoea) · Intra-abdominal infections · Infections of the skin and soft tissue caused by Gram-negative bacteria · Malignant external otitis · Infections of the bones and joints · Inhalation anthrax (post-exposure prophylaxis and curative treatment) 	
K-mentin Duo Syrup Amoxicillin hydrate 36.3143g Potassium clavulanate diluted .. 5.1878g 	Acute and chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, Pelvic infection, gonorrhoea, boils and abscesses, soft tissues, wound infections, osteomyelitis, Dental infection	
Tenofode Tab. Tenofovir Disoproxil Fumarate .. 300mg 	HIV-1 infection, Hepatitis B infection	
Roct Tab. 150mg Roxithromycin 150mg 	<ul style="list-style-type: none"> - acute pharyngitis (sore throat and discomfort when swallowing) - tonsillitis - sinusitis - acute bronchitis (infection of the bronchi causing coughing) - worsening of chronic bronchitis - pneumonia (lung infection characterised by fever, malaise, headache) - skin and soft tissue infections - non gonococcal urethritis - impetigo (bacterial infection causing sores on the skin) 	

KOFERON Inj.

Pegylated interferon alpha-2b

Active Ingredient

Pegylated interferon alpha-2b 120µg

Indications

Chronic hepatitis C

Koferon Inj. is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for hepatitis C virus RNA (HCV-RNA). The best way to use Koferon Inj. in this indication is in combination with ribavirin.

Dosage and administration




Koferon Inj. is administered once a week by subcutaneous injection. Dosage dose varies with combination therapy with ribavirin and monotherapy.

Koferon Inj in combination with ribavirin


Approximately 1.5µg / kg / of Koferon Inj. is administered in combination with a ribavirin capsule. The dose of 1.5µg / kg of Koferon Inj. to be used in combination with ribavirin should be administered using a vial of the appropriate concentration for each weight range. Ribavirin capsules should be taken every morning and evening with food. oncentration for each weight range. Ribavirin capsules should be taken every morning and evening with food.



Antineoplastic & Immunomodulating Agents

Product Name (Active Ingredients & Strength)	Indications	Remark
Koferon Inj. Pegylated Interferon α-2b 120 µg	 Hepatitis C	
Korus Thymomodulin Cap. Thymomodulin 80mg	 Some autoimmune diseases such as various types of severe hepatitis, Lupus erythematosus, Rheumatoid arthritis, Allergic asthma and other immune cells	
Metorex Tab. Methotrexate 2.5mg	 Rheumatoid arthritis, Uncontrolled psoriasis, Acute leukemia, Non-Hodgkin's lymphoma, Soft-tissue and osteogenic sarcomas, Solid tumors particularly breast, lung, head and neck, bladder, cervical, ovarian, and testicular carcinoma	

Antiparasitic Products, Insecticides & Repellents

Product Name (Active Ingredients & Strength)	Indications	Remark
Korus Albendazole Tab. (a)Albendazole 200mg (b)Albendazole 400mg	 Neurocysticercosis, Hydatid Disease	

K-Grel Tab.

Anti-platelet drug to inhibit blood clots in a variety of conditions such as peripheral vascular disease, coronary artery disease, and cerebrovascular disease

Active Ingredient

Clopidogrel Bisulfate 97.875 mg

Indications

Prevention of atherothrombotic events

Adult patients suffering from myocardial infarction (from a few days until less than 35 days), ischaemic stroke (from 7 days until less than 6 months) or established peripheral arterial disease.

Adult patients suffering from acute coronary syndrome

Prevention of atherothrombotic and thromboembolic events in atrial fibrillation.

Dosage and administration

Adults and older people: Clopidogrel should be given as a single daily dose of 75 mg.

Blood & Blood Forming Organs



KOROMON Inj.

Recombinant Human Erythropoietin product

Active Ingredient

Recombinant Human Erythropoietin 2000/4000/10000IU

Indications

- 1) Treatment of anaemia associated with chronic renal failure in patients on dialysis
- 2) Treatment of symptomatic renal anaemia in patients not yet undergoing dialysis
- 3) Prevention of anemia of in infants and adults and in increasing the yield of autologous blood

Dosage and administration

KOROMON injection is for intravenous administration and subcutaneous administration.











KOROMON injection is for intravenous administration and subcutaneous administration.

50 IU/kg body weight, three times a week for 1-2 min by I.V. route.

50 IU/kg body weight, three times a week for 1-2 min by I.V. route.

Blood & Blood Forming Organs



Product Name (Active Ingredients & Strength)	Indications	Remark
Esnase Inj. (a)Streptokinase 750000IU (b)Streptokinase 1500000IU	 Acute myocardial infarction within 12 hours of onset with persistent ST-segment elevation or recent left bundle-branch block	
K-Grel Tab. Clopidogrel Sulfate 97.875mg	 Myocardial infarction (MI), Recent stroke, Established peripheral arterial disease, Acute Coronary Syndrome	
Koromon Prefilled Inj. (a)Recombinant human erythropoietin 2000IU (a)Recombinant human erythropoietin 4000IU	 Symptomatic anemia, Anemia requiring transfusion	
Ciloen Tab. Cilostazol 50mg	 Improvement of ischemic symptoms such as ulcers, pain and cold sensation due to chronic arterial occlusion (Burger's disease, occlusive atherosclerosis, diabetic peripheral vasculopathy, etc.) Suppression of recurrence after cerebral infarction (Except for cardiogenic cerebral embolism)	
Barokan Tab. Cilostazol 100mg Ginkgo biloba Leaf Extract 80mg	 Alternative therapies for combination therapy with cilostazol and Ginkgo biloba Leaf Extract if the effect of cilostazol alone is insufficient for the following diseases: Improvement of ischemic symptoms such as ulcers, pain and cold sensation due to chronic arterial occlusion (Burger's disease, occlusive atherosclerosis, diabetic peripheral vasculopathy, etc.) Suppression of recurrence after cerebral infarction(Except for cardiogenic cerebral embolism)	
Aspharm Enteric Coated Tab. Aspirin 100mg	 <ul style="list-style-type: none"> - Secondary prevention of myocardial infarction. - Prevention of cardiovascular morbidity in patients suffering from stable angina pectoris. - History of unstable angina pectoris, except during the acute phase. - Prevention of graft occlusion after Coronary Artery Bypass Grafting (CABG). - Coronary angioplasty, except during the acute phase. - Secondary prevention of transient ischaemic attacks (TIA) and ischaemic cerebrovascular accidents (CVA), provided intracerebral haemorrhages have been ruled out. - Acute Myocardial infarction 	
Sarpod Tab. Sarpogrelate Hydrochloride ... 100mg	 Improvement of ischemic symptoms such as ulcers, pain and coldness caused by chronic arterial occlusion (Burger's disease, occlusive atherosclerosis, diabetic peripheral vascular disease, etc.)	
Asgrel Cap. Aspirin 100mg Clopidogrel bisulfate 97.875mg	 Use only in patients with clopidogrel and aspirin at the same time. Improvement of atherosclerotic symptoms in patients with acute coronary syndrome or more vascular risk factors, Vitamin K antagonist (VKA) administration is not appropriate, Decreased risk of atherothrombosis and thromboembolism including stroke in patients with atrial fibrillation with low risk of bleeding	
Tigrel Tab. 60mg / 90mg (a)Ticagrelor 60mg (b)Ticagrelor 90mg	 Brilique, co-administered with acetylsalicylic acid (ASA), is indicated for the prevention of atherothrombotic events in adult patients with - acute coronary syndromes (ACS) or - a history of myocardial infarction (MI) and a high risk of developing an atherothrombotic event	
Apixcos Tab. 2.5mg / 5mg (a)Apixaban 2.5mg (b)Apixaban 5.0mg	 Prevention of venous thromboembolic events (VTE) in adult patients who have undergone elective hip or knee replacement surgery. Prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation (NVAF), with one or more risk factors, such as prior stroke or transient ischaemic attack (TIA); age ≥ 75 years; hypertension; diabetes mellitus; symptomatic heart failure (NYHA Class ≥ II). Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults	

Keitoba Tab.

HMG-CoA Reductase Inhibitor

Active Ingredient

Atorvastatin Calcium 10.36 mg(10mg as Atorvatatin)

Indications

- Reduces risk of cardiovascular disease of following patients
 - Adult patients with no clinical evidence of coronary heart disease, but with multiple risk factor of coronary heart disease
 - Type 2 diabetic patients with no clinical evidence of coronary heart disease, but with multiple risk factor of coronary heart disease
 - Adult patients with clinical evidence of coronary heart disease
- Hyperlipidemia
- Supplementary diet therapy that reduces total cholesterol, LDL-cholesterol, Apolipoprotein B for children of ages 10-17 (In case of female, patients after first period) with heterozygous familial hypercholesterolemia despite of dietary therapy)

Dosage and administration

Before receiving or during treatment of atorvastatin, patient should be placed on a standard cholesterol-lowering diet.

Atorvastatin may be given at any time of day with or without food.

- Hyperlipidemia : Dose range of 10-80mg once a day
- Children with heterozygous familial hypercholesterolemia (Age 10-17): 10-20mg once a day
- Patients with renal disorder: There is no need for dose adjustment because renal disease does not affect plasma concentration of atorvastatin or declining effect of LDL-cholesterol
- Elderly : Efficacy and safety of elderly patients older than 70 are similar to those of normal adult patients

Cardiovascular System



Twyn-on Tab. 40/10mg, 40/5mg, 80/5mg

Hypotensive agents

Active Ingredient

Twyn-on Tab. 40/10mg

Telmisartan powder(23.9%) 167.36 (40mg as Telmisartan), Amlodipine besylate 13.87mg (10mg as Amlodipine)

Twyn-on Tab. 40/5mg

Telmisartan powder(23.9%) 167.36 (40mg as Telmisartan), Amlodipine besylate 6.935mg (5mg as Amlodipine)

Twyn-on Tab. 80/5mg

Telmisartan powder(23.9%) 334.72 (80mg as Telmisartan), Amlodipine besylate 16.935mg (5mg as Amlodipine)

Indications

Essential hypertension in which blood pressure is not adequately controlled by amlodipine or telmisartan monotherapy

Dosage and administration

- 40/5mg** : Amlodipine 5mg or telmisartan 40mg is administered to patients whose blood pressure is not adequately controlled.
- 80/5mg** : Amlodipine 5 mg or about 40/5mg administered to patients whose blood pressure is not adequately controlled.
- 40/10mg**: Amlodipine 10 mg is administered to patients whose blood pressure is not adequately controlled.

Cardiovascular System



Twotwoforge Tab. 5mg/80mg, 5mg/160mg, 10mg/160mg

Hypotensive agents

Active Ingredient

Twotwoforge 5mg/80mg film-coated tablets

Amlodipine besylate 6.94mg, Valsartan 80mg

Twotwoforge 5mg/160mg film-coated tablets

Amlodipine besylate 6.94mg, Valsartan 160mg

Twotwoforge 10mg/80mg film-coated tablets

Amlodipine besylate 13.87mg, Valsartan 160mg

Indications

Treatment of essential hypertension.

Twotwoforge is indicated in adults whose blood pressure is not adequately controlled on amlodipine or valsartan monotherapy.

Dosage and administration

The recommended dose of Twotwoforge is one tablet per day.















Twotwoforge 5mg/80mg may be administered in patients whose blood pressure is not adequately controlled with amlodipine 5mg or valsartan 80mg alone.













Twotwoforge 5mg/160mg may be administered in patients whose blood pressure is not adequately controlled with amlodipine 5mg or valsartan 160mg alone.




Twotwoforge 10mg/160mg may be administered in patients whose blood pressure is not adequately controlled with amlodipine 10mg or valsartan 160mg alone or with Twotwoforge 5mg/160mg.

Cardiovascular System









Product Name (Active Ingredients & Strength)	Indications	Remark
Censelia Tab. Titrated Extract of Centella Asiatica 30mg 	Venous Disease: Improvement of symptoms related to veins and lymphatic insufficiency	
Ezit Tab. Ezetimibe 10mg 	Primary Hypercholesterolaemia, Prevention of Cardiovascular Events, Homozygous Familial Hypercholesterolaemia (HoFH)	
Hemorral Tab. Ginkgo Leaf Ext. 14mg Heptaminol Hydrochloride 300mg Troxerutin 300mg 	The following symptoms of varicose veins: Heavy pressure, phlebitis Various symptoms related to hemorrhoids	
K-duet Tab. (a) 5/10mg Amlodipine Besylate 6.94mg Atorvastatin Calcium 10.36mg (a) 5/20mg Amlodipine Besylate 6.94mg Atorvastatin Calcium 20.72mg 	This drug is used only in patients who need to administer both drugs (amlodipine and atorvastatin) at the same time. Hypertension, Chronic stable angina pectoris, Vasospastic (Prinzmetal's) angina Hypercholesterolaemia, Prevention of cardiovascular disease	
Lozatal Plus Tab. Losartan potassium 50mg Hydrochlorothiazide 12.5mg 	Hypertension in which blood pressure is not adequately controlled by monotherapy	
Telmi-on Tab. 40mg/80mg (a) Telmisartan 40mg (b) Telmisartan 80mg 	Hypertension, Cardiovascular prevention	
Telmi-on Plus Tab. 40/12.5mg (a) Telmisartan 40mg (b) Hydrochlorothiazide 12.5mg  80/12.5mg (a) Telmisartan 80mg (b) Hydrochlorothiazide 12.5mg	Treatment of Essential Hypertension	
Kovaltan Tab. 80mg / 160mg (a) Valsartan 80mg (b) Valsartan 160mg 	Hypertension, Recent myocardial infarction, Heart failure	
Amlova Tab. Amlodipine besylate 6.944mg 	Hypertension, Chronic stable angina pectoris, Vasospastic (Prinzmetal's) angina	
Amlova S Tab. 2.5mg S-Amlodipine Besylate Dihydrate 3.69mg Amlova S Tab. 2.5mg S-Amlodipine Besylate Dihydrate 7.38mg 	Myocardial ischemia due to Hypertension, stable angina or angina pectoris	
Ateol Tab. Atenolol 50mg 	Hypertension, Angina pectoris, Cardiac arrhythmias, Myocardial infarction	
K-Tova Tab. 10mg/20mg (a) Atorvastatin Calcium (10mg as Atorvastatin) 10.36mg (a) Atorvastatin Calcium (20mg as Atorvastatin) 21.69mg 	Primary hypercholesterolemia and mixed dyslipidemia, Hypertriglyceridemia, Primary dysbetalipoproteinemia, Homozygous familial hypercholesterolemia, Cardiac valve disease	
Losatal Tab. Losartan Potassium 50mg Losartan Potassium 100mg 	Hypertension Kidney disease in type 2 diabetic patients with hypertension	
Twotwoforge Tab. (a) Valsartan 80mg Amlodipine Besylate 6.94mg (b) Valsartan 160mg Amlodipine Besylate 6.94mg (c) Valsartan 160mg Amlodipine Besylate 13.87mg 	Amlodipine or valsartan monotherapy cannot adequately control blood pressure Essential Hypertension	

Product Name (Active Ingredients & Strength)	Indications	Remark
Korus Mexiletine Cap. (a) Mexiletine Hydrochloride ··· 100mg (b) Mexiletine Hydrochloride ··· 200mg	 (a)  (b) 	Ventricular arrhythmias, Sustained ventricular tachycardia Ventricular arrhythmia Ventricular arrhythmia, Ventricular arrhythmia due to digitalis, contraction of idiopathic ventricle, ventricular arrhythmia in acute myocardial infarction
K-rotin Tab. (a) Rosuvastatin Calcium ····· 5.2mg (b) Rosuvastatin Calcium ····· 10.4mg	 (a)  (b) 	Primary hypercholesterolemia (type IIa), combined hyperlipidemia (type IIb): Dietary supplements Reduced risk for cardiovascular disease
Kallinogen Tab. Kallidinogenase ··········· 50IU		1. Peripheral circulatory disturbances due to the following diseases: hypertension, Meniere's syndrome, obstructive thromboangiitis (Burger's disease) 2. Improvement of various symptoms due to menopausal disorder, circulatory disorder of the mesenteric choroid
Twyn-on Tab. (a) 80/5mg Amlodipine Besylate ······· 6.935mg Telmisartan Powder(23.9%) ··· 334.72mg (b) 40/10mg Amlodipine Besylate ······· 13.87mg Telmisartan Powder(23.9%) ··· 167.36mg (b) 40/10mg Amlodipine Besylate ······· 6.935mg Telmisartan Powder(23.9%) ··· 167.36mg		Essential hypertension; in case of Amlodipine or telmisartan cannot be controlled appropriate
Limart Tab. Limaprost alfadex ········· 166.67µg		1. Improvement of ischemic symptoms such as ulceration, pain caused by thromboangiitis obliterans (Burger's disease) 2. Improvement of subjective symptoms (lower back pain, lower limb) and walking ability by acquired lumbar spinal stenosis (normal, bilateral intermittent claudication in SLR test)
Ezit Tab. Ezetimibe ················· 10mg		Primary Hypercholesterolaemia, Prevention of Cardiovascular Events, Homozygous Familial Hypercholesterolaemia (HoFH)
Hemoral Tab. Ginkgo Leaf Ext. ··········· 14mg Heptaminol Hydrochloride ··· 300mg Troxerutin ················· 300mg		The following symptoms of varicose veins: Heavy pressure, phlebitis Various symptoms related to hemorrhoids
K-simva Tab. 20mg Simvastatin ··············· 20mg		Hypercholesterolaemia, Cardiovascular prevention

Product Name (Active Ingredients & Strength)		Indications	Remark
Muzona Cream Terbinafine hydrochloride 10mg/g		Tinea pedis and tinea cruris caused by Trichophyton (e.g. T. rubrum, T. mentagrophytes, T. verrucosum, T. violaceum) and Epidermophyton floccosum	
Methasone Cream Mometasone furoate 1mg/g		For the treatment of inflammatory and pruritic manifestations of psoriasis (excluding widespread plaque psoriasis) and atopic dermatitis.	
Plimycin Ointment Bacitracin zinc 500IU Neomycin sulfate 3.5mg Polymyxin B sulfate 10000IU Pramoxine HCl 10mg		Prevention of infectious diseases and pain relief due to scab, wounds, burns	

Genitourinary System & Reproductive Hormones

Product Name (Active Ingredients & Strength)		Indications	Remark
Kozosin Tab. Terazosin Hydrochloride Hydrate (1mg as Terazosin) .. 1.187mg/2.374mg		Hypertension (Mild to Moderate), Urinary disorder due to benign prostatic hyperplasia (BPH)	
Flitop Tab. 75mg Naftopidil 75mg		Dysuria due to benign prostatic hyperplasia	
Solfelicin Tab. 5mg/10mg (a) Solifenacin succinate 5mg (b) Solifenacin succinate 10mg	(a)  (b) 	Symptomatic treatment of urge incontinence and/or increased urinary frequency and urgency as may occur in patients with overactive bladder syndrome.	
Provine Tab. 20mg Propiverine Hydrochloride 20mg		Symptomatic treatment of urinary incontinence and/or increased urinary frequency and urgency in patients with overactive bladder syndrome or neurogenic detrusor overactivity from spinal cord injuries	
Finote Tab. Finasteride 5mg		Finote Tab. is indicated for the treatment and control of benign prostatic hyperplasia (BPH) to: - cause regression of the enlarged prostate, improve urinary flow and improve the symptoms associated with BPH, - reduce the incidence of acute urinary retention and reduce need for surgery including transurethral resection of the prostate (TURP) and prostatectomy.	

Hyron Prefilled Inj.

Viscosupplementation agents

Active Ingredient

Sodium Hyaluronate 20mg/2mL, 40mg/2mL

Indications












Hyron Prefilled Inj. is indicated for the treatment of pain in osteoarthritis(OA) of the knee in patients who have failed to respond adequately to conservative non-pharmacologic therapy and simple analgesics, e.g., acetaminophen.










Dosage and administration















Hyron Prefilled Inj. is administered by intra-articular injection once a week (one week apart) for a total of three injections.

Musculoskeletal System



Product Name (Active Ingredients & Strength)	Indications	Remark
Anac Tab. Aceclofenac 100mg	 Pain and inflammation in osteoarthritis, Rheumatoid arthritis and ankylosing spondylitis.	
Cerebrin Cap. Celecoxib 200mg	 Osteoarthritis, Rheumatoid Arthritis, Juvenile Rheumatoid Arthritis, Ankylosing Spondylitis, Acute Pain, Primary Dysmenorrhea	
Ketorolac Tab. (a)Ketorolac Tromethamine 10mg (b)Ketorolac Tromethamine 20mg	 Moderate to severe pain including postoperative pain	
Korus Diclofenac Tab. (a)Diclofenac sodium 25mg (b)Diclofenac sodium 50mg (a)Diclofenac sodium 100mg	 All grades of pain and inflammation in a wide range of conditions, including: i) Arthritic conditions: rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, acute gout ii) Acute musculo-skeletal disorders such as peri-arthritis (for example frozen shoulder), tendinitis, tenosynovitis, bursitis iii) Other painful conditions resulting from trauma, including fracture, low back pain, sprains, strains, dislocations, orthopaedic, dental and other minor surgery.	
Korus Ibuprofen Tab. (a)Ibuprofen 200mg (b)Ibuprofen 400mg	 Mild to moderate pain including rheumatic and muscular pain, Backache, Neuralgia, Migraine, Headache, Dental pain, Dysmenorrhoea, Feverishness and for the relief of the symptoms of cold and influenza	
Korus Ibuprofen Cap. Ibuprofen 200mg		
Korus Mefenamic Acid Tab. (a)Mefenamic acid 250mg (b)Mefenamic acid 500mg	 Rheumatoid arthritis, Osteoarthritis (Degenerative arthritis), Ankylosing spondylitis, Tendinitis, Acute gout, Dysmenorrhea, Bursitis, Musculoskeletal disorder, Postoperative pain, Migraine, Pain after tooth extraction	
Pacisulide Tab. Nimesulide 100mg	 Acute pain (Postoperative pain, Dental pain, Headache, Low back pain)	
Tamate Tab. Talinifumate 370mg	 Rheumatoid arthritis, Osteoarthritis (Degenerative arthritis), Sciatica, Amenorrhic pain, Post-traumatic pain, Postoperative inflammation and pain, Tendovaginitis, Distortion, Uterine adnexitis, Pharyngitis, Tonsillitis, Otitis, Sinusitis	
Tamenbun Tab. Nabumetone 500mg	 Osteoarthritis and rheumatoid arthritis requiring anti-inflammatory and analgesic treatment	
Tamic Cap. Talinifumate 370mg	 Rheumatoid arthritis, Osteoarthritis (Degenerative arthritis), Ankylosing spondylitis, Tendinitis, Acute gout, Dysmenorrhea, Bursitis, Musculoskeletal disorder, Postoperative pain, Migraine, Pain after tooth extraction	

Product Name (Active Ingredients & Strength)	Indications	Remark
Lopenal Tab. Loxoprofen Sodium Hydrate 68.1mg 	1. Symptoms of the following diseases and symptoms Pain Chronic rheumatoid arthritis, osteoarthritis (degenerative arthritis), back pain, shoulder circumflex, cervicitis syndrome 2. post-operative, post-traumatic and post-extraction anti-inflammatory pain 3. Death paralysis of the following diseases Acute topical salts (including acute topical salts with acute bronchitis)	
Erisono Tab. Eperisone Hydrochloride 50mg 	1. Painful muscular spasm associated with musculoskeletal disorders: cervicothoracic syndrome, shoulder joint inflammation, back pain 2. Spinal cord paralysis due to neurological diseases: cerebrovascular disorder, spinal cord paralysis, cervical spondylosis, post-operative sequelae (including brain and spinal cord tumors), trauma sequelae (spinal cord injury, head trauma), amyotrophic sclerosis, cerebral palsy, Degenerative disease, spinal cord vascular disorder, subacute spinal cord optic nerve lesion, other cerebrospinal disease	
Hyron Prefilled Inj. Sodium Hyaluronate 20mg/2mL Sodium Hyaluronate 40mg/2mL 	Osteoarthritis treatment of knee Periarthritis, Gonarthrocace	
Febroct Tab. 40mg / 80mg (a)Febuxostat 40mg (b)Febuxostat 80mg 	Treatment of chronic hyperuricaemia in conditions where urate deposition has already occurred	
Alenstar Tab. Alendronate sodium trihydrate 91.37mg 	Osteoporosis Treatment of postmenopausal women with Men's Osteoporosis Treatment	
Zalto Tab. Zaltoprofen 80mg 	Inflammation and symptoms of chronic rheumatoid arthritis, osteoarthritis, back pain, frozen shoulder, cervical syndrome Post-surgery, post-trauma and post-extraction inflammation and pain	
Diaron Cap. Zaltoprofen 80mg 	Bone (hip & knee joint) arthritis (arthropathy, degenerative arthritis)	
Ibaron Inj. Ibandronate Sodium Monohydrate 3.375mg 	Treatment of osteoporosis in postmenopausal women at increased risk of fracture	
Pirox Gel Piroxicam 5mg/g 	Osteoarthritis and rheumatoid arthritis requiring anti-inflammatory and analgesic treatment	

Product Name (Active Ingredients & Strength)	Indications	Remark
Citicoline Inj (a) Sodium Citicoline (500mg as Citicoline) ····· 522.53mg/2mL (b) Citicoline ··········· 250mg/mL		Decreased consciousness due to head injury or brain surgery, Parkinson's disease, Upper limb function recovery promotion of stroke patient, Acute pancreatitis, Chronic relapsing pancreatitis
Koridol Cap. Sodium Hyaluronate ····· 20mg/2mL Sodium Hyaluronate ····· 40mg/2mL		Osteoarthritis treatment of knee Periarthritis, Gonarthrocase
Korus Acetaminophen Tab. (a) Acetaminophen ········· 300mg (b) Acetaminophen ········· 500mg		Fever and pain due to common cold, Headache, Neuralgia, Myalgia, Dysmenorrhea, Pain after distortion, Toothache, Arthralgia, Rheumatic pain
Ranophen Semi Tab. Acetaminophen ········· 162.5mg Tramadol HCl ··········· 18.75mg		Severity - Severe or chronic pain
Neopezil Tab. (a) Donepezil Hydrochloride ·· 5.215mg (b) Donepezil Hydrochloride ·· 10.43mg		Treatment of Symptoms of Alzheimer's Dementia
Ginkgoex Tab. Ginkgo Biloba Extract ······· 80mg		1. Main effect 1) Treatment of peripheral arterial circulatory disorder (intermittent claudication) 2) dizziness, vascularity and degenerative changes 3) Treatment of dysmorphic brain dysfunction accompanied by dementia such as tinnitus, headache, memory loss, concentration disorder, depression, and dizziness
Neocital Tab. (a) Escitalopram oxalate ····· 6.39mg (b) Escitalopram oxalate ····· 12.77mg		Treatment of major depressive episodes, panic disorder with or without agoraphobia, social anxiety disorder (social phobia), generalised anxiety disorder and obsessive-compulsive disorder.
Gabistar Cap. (a) Gabapentin ··········· 100mg (b) Gabapentin ··········· 300mg		Epilepsy Gabapentin is indicated as adjunctive therapy in the treatment of partial seizures with and without secondary generalization in adults and children aged 6 years and above. Treatment of peripheral neuropathic pain Gabapentin is indicated for the treatment of peripheral neuropathic pain such as painful diabetic neuropathy and post-herpetic neuralgia in adults.
Prometine Tab. Memantine hydrochloride ····· 10mg		Treatment of patients with moderate to severe Alzheimer's disease.
Parkin Cap. 75mg/150mg (a) Pregabalin ············· 75mg (b) Pregabalin ············· 100mg		Neuropathic pain, Epilepsy, Generalised Anxiety Disorder
Rivaslon Cap. 1.5mg / 3.0mg (a) Rivastigmine Hydrogen Tartrat ··························· 2.4mg (b) Rivastigmine Hydrogen Tartrat ··························· 4.8mg		Symptomatic treatment of mild to moderately severe Alzheimer's dementia. Symptomatic treatment of mild to moderately severe dementia in patients with idiopathic Parkinson's disease.
Neocital Tab. 5mg /10mg (a) Escitalopram Oxalate ····· 6.39mg (b) Escitalopram Oxalate ····· 12.77mg		Treatment of major depressive episodes. Treatment of panic disorder with or without agoraphobia. Treatment of social anxiety disorder (social phobia). Treatment of generalised anxiety disorder. Treatment of obsessive-compulsive disorder.
Lotine Cap. 10mg / 20mg (a) Fluoxetine Hydrochloride ·· 11.2mg (b) Fluoxetine Hydrochloride ·· 22.4mg		Depressive disorder, Obsessive-compulsive disorder, Bulimia nervosa, Premenstrual dysphoric disorder
Lotine Cap. 10mg / 20mg (a) Duloxetine hydrochloride ········· 33.7 (30mg as duloxetine) (a) Duloxetine hydrochloride ········· 67.3 (60mg as duloxetine)		Treatment of major depressive disorder. Treatment of diabetic peripheral neuropathic pain. Treatment of generalised anxiety disorder.

Ausmin Cap.

Dietary Supplement

Active Ingredient

DL-Methionine	9.2mg	Tocopherol Acetate 50%	10.0mg
L-Leucine	9.15mg	Thiamine Mononitrate	2.5mg
L-Valine	5.0mg	Riboflavin	1.5mg
L-Isoleucine	5.0mg	Nicotinamide	10.0mg
L-Threonine	5.0mg	Pyridoxine Hydrochloride	5.0mg
L-Tryptophan	5.0mg	Calcium Pantothenate	2.5mg
L-Phenylalanine	5.0mg	Cyanocobalamin	0.5ug
L-Lysine Hydrochloride	25.0mg	Folic Acid	100.0ug
Retinol Palmitate	1,000IU	Ascorbic Acid	25.0mg
Ergocalciferol	100IU		

Indications

Maintenance of physical strength, nutritional supplements for children, aged people, nursing mothers, and convalescent patients.

Dosage and administration

Adults and children over 10 years of age : orally, 1 capsule, 3 times daily.
Children : Orally, 1-2 capsules daily. However, in the case of gastric ulcer, it should be administered before breakfast, evening and bedtime.

Nutrients



Actizin Tab.

Premium Active Vitamin B1 (Dietary Supplement)

Active Ingredient

γ-oryzanol	10mg	Coated Ascorbic acid	52.1mg
Nicotinamide	100mg	Choline Tartrate	20mg
Riboflavin	100mg	Tocopherol Acetate 50%	20mg
Biotin	0.1mg	Calcium Pantothenate	100mg
zinc oxide	30mg	Pyridoxine hydrochloride	100mg
Cyanocobalamin	1mg	Cholecalciferol Concentrated Powder	10mg
Folic acid	50μg	Magnesium oxide	50mg
Ursodeoxycholic acid	10mg	Benfotiamine	100mg
Inositol	100mg	Dried yeast with selenium	20mg

Indications







Prevalence of vitamin D, E, B1, B2, B6, C in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age
The efficacy and effects of vitamins, etc., contained in this medicine are as follows :
poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue, Symptom relief follows: Neuralgia, Myalgia, Arthralgia(back pain, shoulder stiffness, etc.), Angulus infectious, Chelitis, Stomatitis, glossitis, eczema, dermatitis Supplement of zinc




Dosage and administration

Take 1 tablet once a day.


Nutrients/Ophthalmic agent



Product Name (Active Ingredients & Strength)	Indications	Remark
<p>Ausmin Cap.</p> <p>DL-Methionine 9.2mg L-Leucine 9.15mg L-Valine 5mg L-Isoleucine 5mg L-Threonine 5mg L-Tryptophan 5mg L-Phenylalanine 5mg L-Lycine Hydrochloride 25mg Retinol Palmitate 1000IU Ergocalciferol 100IU Tocopherol acetate 50% 10mg Thiamine Nitrate 2.5mg Riboflavin 2.5mg Niacinamide 10mg Pyridoxine Hydrochloride 5mg Calcium Pantothenate 2.5mg Cyanocobalamin 0.5µg Folic Acid 100µg Ascorbic Acid 25mg</p>	 <p>Increased Energy, Fragile habitus, Physical fatigue, Wasting disease, Prevalence of calcium and nutrient during gestational and lactation period</p>	
<p>Magtamine Soft Cap.</p> <p>Tocopherol acetate 500mg Magnesium Oxide 250mg Benfotiamin 20mg Thiamine nitrate 30mg Pyridoxine hydrochloride 50mg γ-Oryzanol 10mg</p>	 <p>Prevalence of vitamin E, B1, B6 in the following cases - Physical fatigue, pregnancy, breastfeeding, illness or illness (during or after illness)</p>	
<p>Double Active Tab.</p> <p>Benfotiamine 25mg Fursultiamine 25mg Riboflavin 50mg Nicotinamide 20mg Calcium Pantothenate 20mg Biotin 0.2mg Folic acid 0.4mg Coated Ascorbic acid 650mg Tocopherol Acetate 50% 20mg zinc oxide 37.35mg Cholecalciferol Concentrated Powder 10mg Ubidecarenone 10mg γ-oryzanol 10mg</p>	 <p>Prevalence of vitamin D, E, B1, B2, C in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue, Relaxation of pigmentation (stain, freckles) caused by sunlight and skin diseases Prevention of Gingival bleeding & nose bleeding Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectiousus, Cheilitis, Stomatitis (mouth sores), glossitis, eczema, dermatitis Supplement of zinc</p>	
<p>Cellbone Vita Tab.</p> <p>γ-oryzanol 10mg Nicotinamide 10mg 0.1 % Cyanocobalamin Powder 20mg Inositol 10mg Choline Tartrate 20mg Calcium Pantothenate 5mg Fursultiamine 50mg Pyridoxine hydrochloride 10mg Sodium Chondroitin Sulfate 600mg Cholecalciferol concentrate powder 10mg Riboflavin Butyrate 50mg</p>	 <p>Prevalence of vitamin B1, B2, B6 in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectiousus, Cheilitis, Stomatitis (mouth sores), glossitis, eczema, dermatitis</p>	
<p>Cen B 100 Tab.</p> <p>γ-oryzanol 5mg Nicotinamide 100mg Riboflavin 100mg Biotin 0.1mg zinc oxide 20mg 0.1 % Cyanocobalamin Powder 20mg Inositol 30mg Tocopherol Acetate 50% 20mg Calcium Pantothenate 100mg Folic acid 0.1mg Pyridoxine hydrochloride 100mg Choline Tartrate 30mg Cholecalciferol Concentrated Powder 10mg Magnesium oxide 50mg Benfotiamine 100mg Dried yeast with selenium 20mg</p>	 <p>Prevalence of vitamin B1, B2, B6 in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectiousus, Cheilitis, Stomatitis (mouth sores), glossitis, eczema, dermatitis</p>	
<p>Benfo B 100 Tab.</p> <p>γ-oryzanol 10mg Nicotinamide 100mg Riboflavin 100mg Biotin 0.1mg zinc oxide 31.2mg 0.1 % Cyanocobalamin Powder 100mg Ursodeoxycholic acid 10mg Ubidecarenone 10mg Inositol 100mg Choline Tartrate 100mg Calcium Pantothenate 100mg Folic acid 0.1mg Pyridoxine hydrochloride 100mg Tocopherol Acetate 50% 100mg Benfotiamine 100mg Magnesium oxide 30mg Coated Ascorbic acid 104.2mg Dried yeast with selenium 9.23mg</p>	 <p>Prevalence of vitamin D, E, B1, B2, B6, C in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue, Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectiousus, Cheilitis, Stomatitis (mouth sores), glossitis, eczema, dermatitis Supplement of zinc</p>	

Product Name (Active Ingredients & Strength)	Indications	Remark
<p>Actizin Tab.</p> <ul style="list-style-type: none"> γ-oryzanol 10mg Nicotinamide 100mg Riboflavin 100mg Biotin 0.1mg zinc oxide 30mg Cyanocobalamine 1mg Folic acid 50µg Ursodeoxycholic acid 10mg Inositol 100mg Coated Ascorbic acid 52.1mg Choline Tartrate 20mg Tocopherol Acetate 50% 20mg Calcium Pantothenate 100mg Pyridoxine hydrochloride 100mg Cholecalciferol concentrate powder .. 10mg Magnesium oxide 50mg Benfotiamine 100mg Dried yeast with selenium 20mg 	 <p>Prevalence of vitamin D, E, B1, B2, B6,C in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age</p> <p>The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue, Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectionsus, Cheilitis, Stomatitis, glossitis, eczema, dermatitis Supplement of zinc</p>	
<p>Megagold Plus Tab.</p> <ul style="list-style-type: none"> γ-oryzanol 5mg Nicotinamide 50mg Riboflavin 6mg Biotin 0.1mg zinc oxide 31.2mg 0.1 % Cyanocobalamine Powder .. 10mg Inositol 25mg Choline Tartrate 20mg Cholecalciferol concentrate powder .. 0.5mg Calcium Pantothenate 15mg Fursultiamine 50mg Pyridoxine hydrochloride 10mg Sodium Chondroitin Sulfate .. 600mg 	 <p>Prevalence of vitamin D, B1, B2, B6 in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age</p> <p>The efficacy and effects of vitamins, etc., contained in this medicine are as follows: Poor development of bones and teeth, prevention of rickets, Beriberi, eye fatigue,</p>	
<p>Neuroelmin Tab.</p> <ul style="list-style-type: none"> Benfotiamine 50mg Hydroxocobalamin Acetate .. 522.5µg Pyridoxal Phosphate Hydrate ... 30mg Riboflavin 10mg Tocopherol Acetate 50% 50mg γ-Oryzanol 5mg 	 <p>Prevalence of vitamin E, B1, B2, B6 in the following cases: Physical fatigue, Pregnancy and lactation, In case of illness or illness (during illness or after recovery), growing, Old age</p> <p>Symptom relief follows: Neuralgia, Myalgia, Arthralgia (back pain, shoulder stiffness, etc.), Angulus infectionsus, Cheilitis, Stomatitis (mouth sores), glossitis, eczema, dermatitis</p>	

■ Ophthalmic Agent

Product Name (Active Ingredients & Strength)	Indications	Remark
<p>Superox Eye Drop</p> <ul style="list-style-type: none"> Ofloxacin 3mg/mL 	 <p>External ocular infections such as conjunctivitis and keratocon junctivitis</p>	

SOMATRON Inj.

Human Growth hormone

Active Ingredient

Somatropin 16IU (5.3 mg)

Indications

Children Growth disturbance due to insufficient secretion of growth hormone (growth hormone deficiency, GHD) and growth disturbance associated with Turner syndrome or chronic renal insufficiency. Adult Growth hormone replacement therapy in adults with growth hormone(GH) deficiency diagnosed is two dynamic tests. Childhood onset & Adult onset

Dosage and administration

The recommended dosage at the start of therapy is 0.125IU (0.04mg) /kg/week given as subcutaneous injection (6-7 divided dose). The dose may be increased gradually according to individual patient requirements to a maximum of 0.25IU (0.08mg)/kg/week.











Occurrence of side effects and serum insulin-like growth factor (IGF-1) levels may be used as guidance for age-and gender-adjusted dose titration.

Minimum effective dose should be used. The dosage should be decreased in cases of persistent edema or severe paresthesia, in order to avoid development of carpal tunnel syndrome.





Systemic Hormonal Preparations








Respiratory System

Product Name (Active Ingredients & Strength)	Indications	Remark
Exos Tab. Triprolidine hydrochloride 2.5mg Pseudoephedrine Hydrochloride .. 60mg	 Allergic rhinitis, Vasomotor rhinitis, Common cold, Influenza	
Korold Tab. Acetaminophen 500mg Chlorpheniramine Maleate 2mg Pseudoephedrine Hydrochloride .. 30mg	 Symptoms of common cold (Runny nose, Nasal obstruction, Sneezing, Throat pain, Chills, Fever, Headache, Arthralgia, Myalgia)	
Betaon Tab. Bepotastine besilat 10mg	 Eczema, dermatitis, skin pruritus,	
Epillergy Tab. (a)Epinastine hydrochloride 10mg (b)Epinastine hydrochloride 20mg	 1. bronchial asthma 2. Allergic rhinitis 3. Normal psoriasis with urticaria, eczema, dermatitis, itchy, itchy rash	
Eloin Cap. Erdosteine 300mg	 Mucus dissolution and expectorants in acute and chronic respiratory diseases	
Monteluk Tab. 10mg Montelukast Sodium 10.4mg	 Prevention and Continuous Treatment of Asthma,. Alleviate seasonal and allergic rhinitis symptoms throughout the year	
Monteluk Chewable Tab. 4mg/5mg (a)Montelukast Sodium 4.16 mg (b)Montelukast Sodium 5.2mg	  Prevention and Continuous Treatment of Asthma,. Alleviate seasonal and allergic rhinitis symptoms throughout the year	
Alti-on Tab. Azelastine Hydrochloride 1mg	 Bronchial asthma, allergic rhinitis, urticaria, eczema, dermatitis, atopic dermatitis, skin pruritus, itching	
Pelisium Tab. Pelargonium Sidaoides 11% Ethanol Extract(1→8-10) 1538.46mg	 Acute bronchitis	







■ Respiratory System

Product Name (Active Ingredients & Strength)	Indications	Remark
Kometasone Nasal Spray Mometasone Furoate 0.5mg/mL		Indicated for use in adults and children 3 years of age and older to treat the symptoms of seasonal allergic or perennial rhinitis. For the treatment of nasal polyps in adults 18 years of age and older.
Korus Levocetirizine Hydrochloride Tab. Levocetirizine Hydrochloride 5mg		The relief of nasal and ocular symptoms of seasonal and perennial allergic rhinitis; The relief of symptoms of chronic idiopathic urticaria.
Mutein Cap. Acetylcysteine 200mg		Difficulty in sputum expectoration in the following diseases: acute, chronic bronchitis, bronchial asthma, laryngitis, sinusitis, cystic fibrosis
Pelisium Syrup Pelargonium sidoides ext. · · 171.6mg/mL		Acute bronchitis

■ Systemic Hormonal Preparations

Product Name (Active Ingredients & Strength)	Indications	Remark
Korus Dexamethasone Inj. Dexamethasone Disodium Phosphate 5mg/mL		Endocrine disease, Rheumatic disease, Collagen disorder, Dermatological disease, Allergic states, Ophthalmological disease, Ulcerative colitis, Blood disease, Leukemia, Nephrotic syndrome, Hodgkin's disease, Urgent case of severe infection
Korus Prednisolone Tab. Prednisolone 5mg		Allergic states, Collagen disorders, Rheumatic disorders, Skin conditions, Neoplastic disease, Gastro-intestinal disease, Respiratory disease, Haematological disorders, Nephrotic syndrome
Somatron Inj. rhGH(Somatropin) 16IU		Growth failure, Growth disturbance, Growth hormone deficiency
Metron Inj. (a)Methylprednisolone Sodium Succinate 29.56mg (b)Methylprednisolone Sodium Succinate 53.0mg (c)Methylprednisolone Sodium Succinate 165.75mg (d)Methylprednisolone Sodium Succinate 331.5mg (e)Methylprednisolone Sodium Succinate 663.0mg	 (a) (b) (c) (d) (e)	Dermatological disease, Allergic states, Gastro-intestinal disease, Respiratory diseases, Neurological disorders, T.B. meningitis, Transplantation
Medisol Tab. Methylprednisolone 4mg		Endocrine disorders, Rheumatic disorders, Collagen diseases/arteritis, Dermatological diseases, Allergic states, Ophthalmic diseases, Respiratory diseases, Haematological disorders, Neoplastic diseases, Gastro-intestinal diseases, Miscellaneous

■ Sensory organs

Product Name (Active Ingredients & Strength)	Indications	Remark
Crolin Eye drops Cromolyn sodium 20mg/mL		Keratitis, conjunctivitis, Allergic conjunctivitis
Korus Sodium Hyaluronate Eye drops Sodium Hyaluronate 1mg/mL		keratoconjunctivitis epithelial defect due to the following diseases: - Endogenous diseases such as Sjogren's syndrome, Stevens Johnson syndrome and dry eye syndrome - Exogenous diseases caused by medication, trauma, wearing contact lenses and post-surgery
K-moxi Eye drops Moxifloxacin hydrochloride · · 5.45mg/mL		Bacterial conjunctivitis, Meibomianitis, Treatment of keratitis (including corneal ulcers), Aseptic therapy during ophthalmic surgery
K-methol Eye drops Fluorometholone 5.45mg/mL		Inflammatory diseases of the Anterior and Posterior segment
Lexacin Eye drops Levofloxacin 5mg/mL		Blepharitis, hordeolum, dacryocystitis, conjunctivitis, keratitis, corneal ulcer, Aseptic therapy during ophthalmic surgery
Oxacin Eye drops Ofloxacin 3mg/mL		Blepharitis, hordeolum, dacryocystitis, conjunctivitis, keratitis, corneal ulcer, Aseptic therapy during ophthalmic surgery